

EFFECTS OF HEAVY MAJORANA NEUTRINOS AT LEPTON-PROTON COLLIDERS

A. Ali^a

Deutsches Elektronen-Synchrotron DESY, Hamburg, Germany

A.V. Borisov^b, D.V. Zhuridov^c

Faculty of Physics, Moscow State University, 119992 Moscow, Russia

Abstract. We discuss the prospects of detecting the processes $e^+p \rightarrow \bar{\nu}_e \ell^+ \ell'^+ X$ and $\nu_e p \rightarrow e \ell^+ \ell'^+ X$ ($\ell, \ell' = e, \mu, \tau$) under the conditions of the present ep collider HERA and of future colliders. These high-energy processes are assumed to be mediated by the exchange of heavy Majorana neutrinos (HMN). We consider two simple scenarios for the HMN mass spectrum: the effective singlet ($m_1 \ll m_2 < m_3 \dots$) and the effective doublet ($m_1 < m_2 \ll m_3 \dots$). For the latter case, the cross section includes information about CP -violating phases.

Introduction

At the moment there are experimental evidences for nonzero neutrino masses [1, 2]. The nature of neutrino mass, whether it is Dirac or Majorana, is one of the fundamental and still unsolved problems in particles physics. A Dirac neutrino carries a lepton number distinguishing a particle from an antiparticle. In contrast to that, a Majorana neutrino is identical to its own antiparticle. The Majorana mass term in the total Lagrangian does not conserve lepton number, but changes its value by two units. Therefore Majorana neutrinos can lead to various lepton number violating processes. For example, they induce same-sign dilepton production in collisions at high energies: $pp \rightarrow \ell^\pm \ell'^\pm X$ [3], $e^+p \rightarrow \bar{\nu}_e \ell^+ \ell'^+ X$ [4] etc.

In theories extending the Standard Model the seesaw mechanism is often used to provide a natural generation of small neutrino masses (for a review, see, e.g., [5, 6]). Unlike the usual way of Dirac mass generation through weak $SU(2)$ -breaking, this mechanism doesn't need extremely small Yukawa couplings ($\lesssim 10^{-12}$). For three families of leptons and s right-handed $SU(2)$ singlets the seesaw mechanism leads to 3 light and s heavy massive Majorana neutrino states

$$\nu_\ell = \sum_{i=1}^3 \tilde{U}_{\ell i} \nu_i + \sum_{j=1}^s U_{\ell j} N_j,$$

where ν_ℓ is a neutrino of definite flavor ($\ell = e, \mu, \tau$), the coefficients $\tilde{U}_{\ell i}$ and $U_{\ell j}$ form the leptonic mixing matrices.

Heavy mass states give a relatively small contribution to neutrino flavor states. Nevertheless effects of light and heavy Majorana neutrinos (HMN)

^ae-mail: ahmed.ali@desy.de

^be-mail: borisov@ave.phys.msu.su

^ce-mail: jouridov@mail.ru

compete in lepton number violating processes, because small values of the mixing parameters $U_{\ell j}$ for heavy neutrinos N_j may be compensated by smallness of the masses of light neutrinos ν_i .

The process $e^+p \rightarrow \bar{\nu}_e \ell^+ \ell'^+ X$

In this report, we investigate the possibilities of observation of the process

$$e^+p \rightarrow \bar{\nu}_e \ell^+ \ell'^+ X \quad (1)$$

and its cross symmetric process $\nu_e p \rightarrow e \ell^+ \ell'^+ X$ (X denotes hadron jets) under the conditions of the present ep collider HERA (DESY) [1] and of future ep colliders. We assume that these processes at high energies

$$\sqrt{s} \gg m_W$$

are mediated by HMN. The leading-order Feynman diagram for the process (1) is shown in Fig. 1. (There is also a crossed diagram with interchanged lepton lines.) For calculating the cross sections, we use the leading effective vector-boson (EVB) approximation [7] neglecting transverse polarizations of W bosons and quark mixing. For this case, cross sections for the process and the crossed channel turn to be equal. As an observation criteria for the process we have chosen the condition

$$\sigma L \geq 1,$$

where σ denotes the cross section and L is the integrated luminosity per year for a collider.




Figure 1: Feynman diagram for the process $e^+p \rightarrow \bar{\nu}_e \ell^+ \ell'^+ X$ mediated by the HMN, N .

We should note that lepton-proton collisions are free of the Standard Model background [4] in contrast to the proton-proton collisions [8].

Effective Singlet Case

At first we take the simplest pattern of the HMN mass spectrum

$$m_1 \ll m_2 < m_3 \dots$$

($m_{N_i} \equiv m_i$) assuming the condition to be held

$$\sqrt{s} \ll m_2.$$

The cross section

$$\sigma_1 = C \left(1 - \frac{1}{2}\delta_{\ell\ell'}\right) |U_{\ell 1} U_{\ell' 1}|^2 \left(\frac{m_1}{m_W}\right)^2 \int_{y_0}^1 \frac{dy}{y} \int_y^1 \frac{dx}{x} p(x, xs) h\left(\frac{y}{x}\right) \omega\left(\frac{ys}{m_1^2}\right) \quad (2)$$

for the process is determined by the convolution of three functions given in [3]: $p(x, xs)$, the quark distribution density having a fraction x of the proton momentum evaluated at the scale $Q^2 = xs$, h , the normalized luminosity of W^+W^+ pairs in the quark-lepton system, and ω , the normalized cross section for the subprocess $W^+W^+ \rightarrow \ell^+\ell'^+$. Here, $y_0 = 4m_W^2/s$ and the characteristic constant C has the value

$$C = G_F^4 m_W^6 / (8\pi^5) = 0.80 \text{ fb}. \quad (3)$$

In the numerical calculation we have used the set of parton distributions CTEQ6 [9]. Using the bounds on the mixing parameters $U_{\ell N}$ from precision electroweak data [10]

$$\sum_N |U_{eN}|^2 < 6.6 \times 10^{-3}, \sum_N |U_{\mu N}|^2 < 6.0 \times 10^{-3}, \sum_N |U_{\tau N}|_{eff}^2 < 3.1 \times 10^{-3} \quad (4)$$

and the constraint from the neutrinoless double beta decay [11]

$$\left| \sum_N U_{eN}^2 m_N^{-1} \right| < 5 \times 10^{-5} \text{ TeV}^{-1}$$

(the sum is over the heavy neutrinos), we find that the process is practically unobservable at HERA even with a very optimistic luminosity ($\sqrt{s} = 318 \text{ GeV}$, $L = 1 \text{ fb}^{-1}$) and also at the projected supercollider VLHC (see, e.g., [12]) ($\sqrt{s} = 6320 \text{ GeV}$, $L = 1.4 \text{ fb}^{-1}$). For example, for $m_1 \sim 1 \text{ TeV}$, we get $\sigma L \sim 10^{-10}$ (10^{-3}) for HERA (VLHC). For a possible detection of the process, the luminosity and/or the energy of the ep -collider should be substantially increased. Taking for example the luminosity $L = 100 \text{ fb}^{-1}$, the observation of the most probable events ($\mu\tau$ and $\mu\mu$) is possible if $\sqrt{s} > 23 \text{ TeV}$. For $\sqrt{s} = 25 \text{ TeV}$ such a collider will be sensitive to a range of neutrino masses about 1–3 TeV.

Effective Doublet Case

We consider also the neutrino mass spectrum of the effective doublet type

$$m_1 < m_2 \ll m_3 \cdots$$

with the bound on energy $\sqrt{s} \ll m_3$. In this scenario, the cross section for the process (1)

$$\sigma_2 = \frac{C}{2} \int_{y_0}^1 \frac{dy}{y} \int_y^1 \frac{dx}{x} p(x, xs) h\left(\frac{y}{x}\right) W\left(\frac{ys}{m_1^2}, \frac{ys}{m_2^2}\right) \quad (5)$$

includes the normalized cross section for the subprocess

$$W(t_1, t_2) = m_W^{-2} [\rho_1^2 m_1^2 \omega(t_1) + 2c\rho_1\rho_2 m_1 m_2 \Omega(t_1, t_2) + \rho_2^2 m_2^2 \omega(t_2)], \quad (6)$$

which contains the individual contributions of the neutrinos N_1 and N_2 , $\omega(t_i)$, and the interference of the two, $\Omega(t_1, t_2)$, where

$$\Omega(t_1, t_2) = 2 - \frac{1}{t_1 + t_2 + t_1 t_2} \left[\frac{t_2(t_1^2 - 2t_1 t_2 - 2t_2)}{t_1(t_1 - t_2)} \ln(1 + t_1) + (t_1 \leftrightarrow t_2) \right];$$

$$\omega(t) = \lim_{t' \rightarrow t} \Omega(t, t') = 2 + \frac{1}{1+t} - \frac{2(3+2t)}{t(2+t)} \ln(1+t).$$

The mixing parameters for different $\ell\ell'$ channels of the process are

$$\rho_i = \sqrt{2 - \delta_{\ell\ell'}} |U_{\ell i} U_{\ell' i}|, \quad c = \cos \delta_{\ell\ell'}$$

with

$$\delta_{\ell\ell'} = \phi_1 - \phi_2 \in [0, 2\pi), \quad \phi_i = \arg(U_{\ell i} U_{\ell' i}).$$

The phases $\delta_{\ell\ell'}$ carry information about CP-violation. We assume the satu-




Figure 2: The dependence of σ_2 (in attobarn) on m_2 (in TeV) plotted for $r = 0, 1/4, 1/2, 3/4, 1$ with $\sqrt{s} = 25$ TeV, $m_1 = 1.3$ TeV and $c = 1$. The horizontal line DL is the discovery limit.

ration of the upper bound $B = 6.0 \times 10^{-3}$ in the second sum in (4) only by the first two terms, i.e., $|U_{\mu 1}|^2 = rB$, $|U_{\mu 2}|^2 = (1-r)B$ with $r \in [0, 1]$. Then for the most probable $\mu\mu$ channel we obtain

$$\sigma_2 = A(r^2 f_1 + 2cr\bar{r}F_{12} + \bar{r}^2 f_2), \quad \bar{r} = 1 - r, \quad (7)$$

where $f_i = f(s, m_i)$ and $F_{ij} = F(s, m_i, m_j)$ are expressed through obvious convolutions of the functions ω and Ω with h and p , respectively (see Eq. (5)), the constant A has the value $A = 1.4 \times 10^{-5}$ fb. For $r = 1$ ($r = 0$), only a single neutrino N_1 (N_2) contributes to the cross section which is reduced to the form given in Eq. (2). Generalization to the case of n neutrinos is straightforward.

In our calculations, we have chosen the following values for the parameters: $\sqrt{s} = 25$ TeV, $m_1 = 1.3$ TeV, $c = 1$. The cross section (7) as a function of m_2 for various fixed values of r is shown in Fig. 2. For the almost degenerate doublet ($m_1 \simeq m_2$) case and/or for the case of small mixing with N_2 ($r \simeq 1$), the cross section σ_2 is close to σ_1 , the cross-section for the effective singlet case. But we should note that for the case of destructive interference of the two almost degenerate massive states (e.g., for $m_2 \simeq m_1$, $r = 1/2$, $c = -1$), the cross section is vanishingly small.

Acknowledgments

We thank Dmitri Peregoudov for help in numerical calculations.

References

- [1] Particle Data Group Collab.: K. Hagiwara et al., *Phys. Rev.* **D66**, 010001 (2002).
- [2] C. Giunti, E-print Archive: hep-ph/0310238;
A.Yu. Smirnov, E-print Archive: hep-ph/0311259.
- [3] A. Ali, A.V. Borisov, N.B. Zamorin, in “*Frontiers of Particle Physics*” (Proceedings of the 10th Lomonosov Conference on Elementary Particle Physics), ed. by A. Studenikin (World Scientific, Singapore, 2003) p. 74; *Eur. Phys. J.* **C21**, 123 (2001) [hep-ph/0104123].
- [4] M. Flanz, W. Rodejohann, K. Zuber, *Phys. Lett.* **B473**, 324 (2000);
W. Rodejohann, K. Zuber, *Phys. Rev.* **D62**, 094017 (2000).
- [5] P. Langacker, *Nucl. Phys. B (Proc. Suppl.)* **100**, 383 (2001).
- [6] B. Kayser, E-print Archive: hep-ph/0211134.
- [7] S. Dawson, *Nucl. Phys.* **B249**, 42 (1985);
I. Kuss, H. Spiesberger, *Phys. Rev.* **D53**, 6078 (1996).
- [8] A. Datta, M. Guchait, D.P. Roy, *Phys. Rev.* **D47**, 961 (1993).
- [9] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P. Nadolsky, W.K. Tung, E-print Archive: hep-ph/0201195.
- [10] E. Nardi, E. Roulet, D. Tommasini, *Phys. Lett.* **B344**, 225 (1995).
- [11] G. Belanger, F. Boudjema, D. London, H. Nadeau, *Phys. Rev.* **D53**, 6292 (1996).
- [12] M. Blaskiewicz et al., Fermilab Report TM-2158, 29 June 2001;
F.M.L. de Almeida Jr., Y.A. Coutinho, J.A. Martins Simões, M.A.B. do Vale, *Phys. Rev.* **D65**, 115010 (2002).